

Topic 13 -

Euler Method



Sometimes you don't have a method to find an exact solution to a differential equation. So instead you approximate the solution.

One method to do this is called Euler's method.

Its used for first-order initial-value problems:


$$y' = f(x, y)$$

$$y(x_0) = y_0$$

The idea goes like this: Suppose you know that a function y satisfies $y(a) = b$ and $y'(a) = m$.

Then suppose you want to approximate $y(a+h)$ where $h > 0$ is a small number.

If you go h distance along the x -axis and follow the tangent line then the y -value will be $y = b + hm$.



You just iterate this over and over to get your approximation.

Let's use this idea in an example.

Consider the initial-value problem

$$\begin{aligned} y' &= xy \\ y(0) &= 1 \end{aligned}$$

$$\begin{aligned} y' - xy &= 0 \\ e^{-\frac{1}{2}x^2} y' - x e^{-\frac{1}{2}x^2} y &= 0 \end{aligned}$$

$$(y e^{-\frac{1}{2}x^2})' = 0$$

$$y e^{-\frac{1}{2}x^2} = C$$

$$y = C e^{\frac{1}{2}x^2}$$

$$y(0) = 1 \rightarrow C = 1$$

$$y = e^{\frac{1}{2}x^2}$$

From earlier methods we know the solution is $y = e^{\frac{1}{2}x^2}$.


Let's pretend that we don't know this.

Let's try to approximate the solution when $0 \leq x \leq 1$.

First let's divide up $0 \leq x \leq 1$ into smaller segments.

$$\text{Let } h = 0.25 = \frac{1-0}{4}.$$

Using h we can



break $0 \leq x \leq 1$ into 4 equally sized segments. We will approximate the solution to $y' = xy$, $y(0) = 1$ at these four points.

The formula $y' = xy$ tells us the slope of the tangent line of the solution at any point. We can use this to approximate the solution $y = e^{\frac{1}{2}x^2}$ without knowing the solution.

Use the initial-value $y(0) = 1$ to get the first point in our approximation.


Let $x_0 = 0$, $y_0 = 1$.

The tangent line has slope

$$y' = x_0 y_0 = 0$$

at this point.

Move $h = 0.25$ along the tangent line to get the next approximate



point (x_1, y_1) . This is:

$$x_1 = x_0 + h = 0 + 0.25 = 0.25$$

$$y_1 = y_0 + h \cdot \underbrace{y'(x_0, y_0)}_{x_0, y_0}$$

$$= 1 + 0.25(0)(1)$$

$$= 1$$

Now do this again but at $(x_1, y_1) = (0.25, 1)$

The tangent line at $(0.25, 1)$ has slope

$$y'(0.25, 1) = (0.25)(1) = 0.25$$


$$\boxed{y' = xy}$$

Set the next approximate point to be

$$x_2 = x_1 + h = 0.25 + 0.25 = 0.5$$

$$y_2 = y_1 + h \cdot y'(x_1, y_1)$$

$$= 1 + 0.25(0.25)(1) = 1.0625$$




Keep going...

$$x_3 = x_2 + h = 0.5 + 0.25 = 0.75$$

$$y_3 = y_2 + h y'(x_2, y_2)$$

$$= 1.0625 + 0.25 \underbrace{(0.5)(1.0625)}_{y' = 0.53125}$$

$$= 1.19531$$




Last point...

$$x_4 = x_3 + h = 0.75 + 0.25 = 1$$

$$y_4 = y_3 + h y'(x_3, y_3)$$

$$= 1.19531 + 0.25 \underbrace{(0.75)(1.19531)}_{y' = 0.89643}$$

$$= 1.41943$$



Summary

n	x_n	y_n	actual solution $y = e^{\frac{1}{2}x^2}$ evaluated at x_n
0	0	1	1
1	0.25	1	1.03174
2	0.5	1.0625	1.13315
3	0.75	1.19531	1.32478
4	1	1.41943	1.64872

If we used a smaller h we would get a way better approximation. This example is to give an idea of how the method works.

Euler's method

Suppose we want to approximate a solution to

$$y' = f(x, y)$$

$$y(x_0) = y_0$$

Pick some $h > 0$.

We are given x_0, y_0 above.

Set

$$x_n = x_{n-1} + h$$

$$y_n = y_{n-1} + h \cdot f(x_{n-1}, y_{n-1})$$

for $n \geq 1$

Ex: Approximate a solution to

$$\begin{cases} y' = xy \\ y(0) = 1 \end{cases}$$

$$\begin{cases} f(x,y) = xy \\ x_0 = 0, y_0 = 1 \end{cases}$$

on the interval $0 \leq x \leq 0.5$ using $h = 0.1$

The Euler equations here are

$$\begin{cases} x_n = x_{n-1} + h \\ y_n = h \cdot f(x_{n-1}, y_{n-1}) \end{cases}$$

$$\begin{cases} x_n = x_{n-1} + 0.1 \\ y_n = (0.1) x_{n-1} y_{n-1} \end{cases} \quad \left. \vphantom{\begin{cases} x_n = x_{n-1} + 0.1 \\ y_n = (0.1) x_{n-1} y_{n-1} \end{cases}} \right\} n \geq 1$$
$$\begin{cases} x_0 = 0 \\ y_0 = 1 \end{cases}$$

For $n=1$:

$$\begin{aligned} x_1 &= x_0 + h = 0 + 0.1 = 0.1 \\ y_1 &= y_0 + h \cdot x_0 \cdot y_0 \\ &= 1 + (0.1)(0)(1) \\ &= 1 \end{aligned}$$

$$\begin{cases} x_1 = 0.1 \\ y_1 = 1 \end{cases}$$

$$x_2 = x_1 + h = 0.1 + 0.1 = 0.2$$

$$y_2 = y_1 + h \cdot x_1 \cdot y_1$$

$$= 1 + (0.1)(0.1)(1)$$

$$= 1.01$$

$$x_2 = 0.2$$

$$y_2 = 1.01$$

$$x_3 = x_2 + h = 0.2 + 0.1 = 0.3$$

$$y_3 = y_2 + h \cdot x_2 \cdot y_2$$

$$= 1.01 + (0.1)(0.2)(1.01)$$

$$= 1.0302$$

$$x_3 = 0.3$$

$$y_3 = 1.0302$$

$$x_4 = x_3 + h = 0.4$$

$$y_4 = y_3 + h \cdot x_3 \cdot y_3$$

$$= 1.0302 + (0.1)(0.3)(1.0302)$$

$$= 1.061106$$

$$x_4 = 0.4$$

$$y_4 = 1.061106$$

$$x_5 = x_4 + h = 0.5$$

$$y_5 = y_4 + h \cdot x_4 \cdot y_4$$

$$= 1.061106 + (0.1)(0.4)(1.061106)$$

$$= 1.10355024$$

$$x_5 = 0.5$$

$$y_5 = 1.10355024$$

x_n	y_n	actual value of solution $e^{\frac{1}{2}x^2}$ at x_n	approximation we initially did with $h=0.25$
0	1	1	1
0.1	1	1.00501	
0.2	1.01	1.0202	
0.3	1.0302	1.04603	
0.4	1.06106	1.08329	
0.5	1.10355024	1.13315	1.0625

$h=0.25$

1